

A complete sample of twelve very X-ray luminous galaxy clusters at $z > 0.5$ H. Ebeling¹, E. Barrett¹, D. Donovan¹, C.-J. Ma¹, A.C. Edge², L. van Speybroeck³**ABSTRACT**

We present the statistically complete and cosmologically most relevant subset of the twelve most distant galaxy clusters detected at $z > 0.5$ by the MAssive Cluster Survey (MACS). Ten of these systems are new discoveries; only two (MACSJ0018.5+1626 aka CL 0016+1609, and MACSJ0454.1–0300 aka MS 0451.6–0305) were previously known. We provide fundamental cluster properties derived from our optical and X-ray follow-up observations as well as the selection function in tabulated form to facilitate cosmological studies using this sample.

Subject headings: galaxies: clusters: general — cosmology: observations — X-rays: general

1. Introduction

The Massive Cluster Survey (MACS) was launched in 1998 with the goal of compiling the first large, statistical sample of very X-ray luminous clusters at $z > 0.3$. A comparison of the flux limits and solid angles of various X-ray cluster surveys conducted previously demonstrates that the ROSAT All-Sky Survey (RASS) was then – and remains until today – the only existing X-ray dataset that allows these extremely rare systems to be detected in significant numbers (fig. 1 of Ebeling, Edge & Henry 2001). A detailed description of the MACS survey strategy is given in the same paper. In brief, the MACS cluster sample is compiled by applying limits in X-ray flux and spectral hardness to all sources listed in the RASS Bright Source Catalogue (BSC, Voges et al. 1999) that fall within the part of the extragalactic sky that is observable from Mauna Kea ($-40^\circ > \delta > 80^\circ$, $|b| > 20^\circ$) and by conducting an identification program that combines searching various astronomical object catalogues with extensive optical imaging observations. Likely cluster candidates are then targeted in spectroscopic observations and added to the MACS catalogue if their measured redshift exceeds $z = 0.3$. Although the required optical follow-up observations are currently complete only for the MACS Bright Sample (defined as comprising all sources with BSC detect fluxes in excess of 2×10^{-12} erg⁻¹ cm⁻² in the 0.1–2.4 keV band, Ebeling et al. 2007, in preparation), we gave priority to the most distant cluster candidates regardless of X-ray flux down to the overall MACS flux limit of 1×10^{-12} erg⁻¹

¹Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

²Department of Physics, University of Durham, South Road, Durham, DH1 3LE, UK

³Harvard-Smithsonian Center for Astrophysics, 60 Garden St, Cambridge, MA 02138, USA

cm^{-2} . We are thus confident that the sample presented here, the set of the twelve MACS clusters at $z > 0.5$, is complete at the nominal 90% level aimed at for the entire survey.

We assume a Λ CDM cosmology with $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$ and $H_o = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ throughout.

2. Follow-up observations

As they were detected in the course of the survey, MACS clusters at $z > 0.5$ were given highest priority in our in-depth follow-up observations. Eight of the twelve clusters forming the final sample were targeted in dedicated observations at 30 GHz with the BIMA interferometer; a strong detection of the Sunyaev-Zeldovich effect was obtained for all of them (LaRoque et al. 2003). All newly discovered clusters were observed with the ACIS-I instrument aboard the Chandra X-ray Observatory in guaranteed time made available by Chandra Telescope Scientist Leon van Spectroscopy. In December 2006 we completed moderately deep observations with the SuprimeCam wide-field imager on the Subaru 8.3m telescope on Mauna Kea in the B, V, R, I, and z' passbands. In addition, U-band images of all systems have been obtained with the MegaCam 1-degree imager on the 3.6m Canada-France-Hawaii Telescope. High-resolution imaging of the cluster cores was performed with the Advanced Camera for Surveys on the Hubble Space Telescope (Program ID 09722, PI Ebeling) in the F555W and F814W filters. Returning to groundbased observations, we conducted extensive spectroscopic observations of galaxies in the fields of all twelve clusters with multi-object spectrographs on 8m-class telescopes on Mauna Kea (Gemini, Keck-I, Keck-II); a catalogue of more than 3,000 redshifts of galaxies in MACS cluster fields can be found in Barrett & Ebeling (2007).

A wealth of science results, ranging from an in-depth study of the massive cooling-core cluster MACSJ1423.8+2404, through work on galaxy evolution in clusters, to a study of the large-scale structure surrounding the most distant MACS clusters, has been obtained from these observations (LaRoque et al. 2003, Ebeling et al. 2004, Allen et al. 2004, Stott et al. 2007, Smith et al. 2007, Barrett et al. 2007, Kartaltepe et al. 2007, Ebeling et al. 2007, Edge et al. 2007) with additional major projects such as a comprehensive weak-lensing analysis of the cluster mass distribution and two complementary studies yielding strong cosmological constraints on dark matter and dark energy using MACS being well advanced (Donovan et al., in preparation; Mantz et al., in preparation; Allen et al., in preparation).

3. The twelve most distant MACS clusters

To facilitate the use of this statistically complete sample in studies conducted by the scientific community in general we here present the MACS $z > 0.5$ catalogue listing the clusters' fundamental physical properties, such as X-ray position, redshift, radial velocity dispersion, X-ray fluxes and luminosities, and X-ray gas temperatures (Table 1). To allow the computation of search volumes for cosmological and astrophysical applications we also list the X-ray detect flux from the RASS BSC and provide a look-up table of the MACS selection function in Table 2.

Color images of the central 10×10 arcmin² of each cluster field (corresponding to roughly 3.8 Mpc at the median redshift of the clusters in our sample) are shown in Fig. 1. All images are based on VRz/ imaging obtained with SuprimeCam in sub-arcsec seeing. Overlaid are isodensity contours of the adaptively smoothed X-ray emission in the 0.5 – 7 keV band as observed with Chandra. Note the wide range of X-ray morphologies, indicating that the MACS sample is neither biased in favor of cooling-core systems (which feature the highest central X-ray surface brightness levels) nor in favor of heavily disturbed clusters (which may be temporarily boosted in their X-ray luminosity and temperature due to merger-induced shocks). A qualitative assessment of the morphology of the cluster gas as traced by the X-ray surface brightness distribution is given in the final column of Table 1.

4. Summary

We present the statistically complete sample of the twelve most distant MACS clusters, all of which feature redshifts of $z > 0.5$ and ten of which are new discoveries. Extensive follow-up observations at wavelengths from the X-ray to the radio passbands have confirmed that these are indeed exceptionally massive systems and thus the high-redshift counterparts of the best studied massive clusters in the local universe. Results from several studies on individual clusters or the sample as a whole have been published; additional studies, ranging from a comprehensive weak-lensing analysis to cosmological applications of the MACS sample, are well advanced. In recognition of the legacy character of this sample we provide in this paper an overview of fundamental properties of all twelve clusters as well as a tabulated version of the MACS selection function to facilitate the use of our sample by the scientific community for a wide range of astrophysical and cosmological applications.

We dedicate this paper to the memory of our friend and colleague Leon van Speybroeck whose advice and enthusiasm (not to mention his Guaranteed Time on Chandra) inspired all of us. We miss you, Leon. HE acknowledges financial support from NASA and SAO (grants NAG 5-8253 and GO2-3168X, respectively).

REFERENCES

- Allen S.W., Schmidt R.W, Ebeling H., Fabian A.C., van Speybroeck L., 2004, MNRAS, 353, 457
Barrett E. & Ebeling H. 2007, MNRAS, submitted
Barrett E., Ebeling H. & Ma C.-J., 2007, MNRAS, submitted
Ebeling H., Edge A.C. & Henry J.P. 2001, ApJ, 553, 668
Ebeling H., Barrett E. & Donovan D., 2004, ApJ, 609, 49L
Ebeling H., White D.A. & Rangarajan F.V.N. 2006, MNRAS, 368, 65
Ebeling H. et al. 2007, MNRAS, submitted
Edge A.C. et al. 2007, MNRAS, in press

Kartaltepe J., Ebeling H., Donovan D., Ma C.-J. 2007, MNRAS, submitted
 LaRoque S. et al. 2003, ApJ, 583, 559
 Smith G. et al. 2007, MNRAS, submitted
 Stott J. et al. 2007, ApJ, in press
 Voges, W. et al. 1999, A&AS, 349, 389

Table 1. Fundamental cluster properties for the MACS $z > 0.5$ sample.

MACS name	α (J2000)	δ (J2000)	z	n_z	σ (km/s)	$f_{X,\text{det,BSC}}$	$L_{X,\text{BSC}}$	$f_{X,\text{Chandra}}$	$L_{X,\text{Chandra}}$	kT (keV)
MACSJ0018.5+1626	00 18 33.835	+16 26 16.64	0.5461	51	1420	1.33 ± 0.36	14.2 ± 3.9	2.14 ± 0.03	19.6 ± 0.3	9.4 ± 1.1
MACSJ0025.4–1222	00 25 29.381	–12 22 37.06	0.5843	31	730	1.10 ± 0.26	13.7 ± 3.3	0.81 ± 0.02	8.8 ± 0.2	7.1 ± 0.8
MACSJ0257.1–2325	02 57 09.151	–23 26 05.83	0.5053	24	900	1.23 ± 0.28	11.3 ± 2.6	1.80 ± 0.03	13.7 ± 0.3	10.5 ± 1.1
MACSJ0454.1–0300	04 54 11.125	–03 00 53.77	0.5377	27	1250	1.55 ± 0.32	17.1 ± 3.5	1.88 ± 0.04	16.8 ± 0.6	7.5 ± 0.7
MACSJ0647.7+7015	06 47 50.469	+70 14 54.95	0.5908	37	900	1.34 ± 0.29	16.8 ± 3.6	1.49 ± 0.03	15.9 ± 0.4	11.5 ± 1.1
MACSJ0717.5+3745	07 17 30.927	+37 45 29.74	0.5460	139	1640	1.92 ± 0.32	20.9 ± 3.5	2.74 ± 0.03	24.6 ± 0.3	11.6 ± 0.8
MACSJ0744.8+3927	07 44 52.470	+39 27 27.34	0.6972	40	1050	1.19 ± 0.26	21.0 ± 4.6	1.44 ± 0.03	22.9 ± 0.6	8.1 ± 0.7
MACSJ0911.2+1746	09 11 11.277	+17 46 31.94	0.5049	26	1150	1.05 ± 0.26	9.8 ± 2.5	1.00 ± 0.02	7.8 ± 0.3	8.8 ± 0.8
MACSJ1149.5+2223	11 49 35.093	+22 24 10.94	0.5445	64	1810	1.20 ± 0.24	13.0 ± 2.6	1.95 ± 0.04	17.6 ± 0.4	9.1 ± 0.8
MACSJ1423.8+2404	14 23 47.663	+24 04 40.14	0.5428	45	1310	1.03 ± 0.23	11.3 ± 2.5	1.80 ± 0.06	16.5 ± 0.7	7.0 ± 0.7
MACSJ2129.4–0741	21 29 26.214	–07 41 26.22	0.5891	47	1460	1.05 ± 0.28	12.6 ± 3.3	1.45 ± 0.03	15.7 ± 0.4	8.1 ± 0.7
MACSJ2214.9–1359	22 14 57.415	–14 00 10.78	0.5026	65	1290	1.42 ± 0.33	12.6 ± 2.9	1.85 ± 0.03	14.1 ± 0.3	8.8 ± 0.8

Note. — The listed X-ray centroids are determined from Chandra ACIS-I data; redshifts, velocity dispersions and number of redshifts are determined within a circle of 1 arcmin around the X-ray centroid. X-ray fluxes (*detect fluxes* in units of 10^{-12} erg s $^{-1}$ cm $^{-2}$) and luminosities (*total luminosities* in units of 10^{44} erg s $^{-1}$) are for the 0.1–2.4 keV band. MACSJ0025.4–1222 and MACSJ0454.1–0300 are rediscoveries of CL 0016+1609 and MS 0451.6–0305, respectively. Fluxes and luminosities from Chandra exclude X-ray point sources associated to a radius of r_{200} . X-ray temperatures were measured from Chandra data within r_{1000} but excluding a central region of 70 kpc radius around the listed X-ray centroid. Cluster morphology is assessed visually based on the amount of substructure apparent in the X-ray contours as well as overall elongation or sphericity – the assigned codes range from 1 (appearing normal) to 5 (extremely disturbed).

Table 2. MACS selection function.

$f_{\text{det,BSC}}$	solid angle	$f_{\text{det,BSC}}$	solid angle
0.550	0	0.925	10136
0.575	20	0.950	11328
0.600	94	0.975	12501
0.625	251	1.000	13544
0.650	667	1.100	16257
0.675	1253	1.200	17823
0.700	2071	1.300	18909
0.725	2877	1.400	19612
0.750	3882	1.500	20043
0.775	4708	2.000	21123
0.800	5464	3.000	21886
0.825	6205	5.000	22297
0.850	7020	10.000	22488
0.875	7940	50.000	22617
0.900	8999		

Note. — RASS detect fluxes $f_{\text{det,BSC}}$ in units of 10^{-12} erg s $^{-1}$ cm $^{-2}$ in the 0.1–2.4 keV band and solid angle in square degrees covered at fluxes exceeding $f_{\text{det,BSC}}$.

Fig. 1.— VRz/ color images of the twelve MACS clusters at $z > 0.5$ as obtained with the Subaru Suprime-Cam wide-field imager. Exposure times range from 20 to 45 minutes in the three passbands. Only the central 10×10 arcmin² region is shown. Overlaid are logarithmically spaced isodensity contours of the adaptively smoothed X-ray surface brightness in the 0.5–7 keV band as observed with Chandra’s ACIS-I detector. We use the ASMOOTH algorithm of Ebeling, White & Rangarajan (2006) and require a minimal significance of 3σ for all features in the adaptively smoothed X-ray image. (files too large for astroph – all images available from first author upon request)